

S/181/61/003/005/008/042
B101/B214

Derivation of a kinetic ...

This equation can be expressed by Green's temperature functions of the complex time $t + i\beta'$. A different method is, however, followed in the following: When use is made of the cyclic property of the spur and the identity:

$[g, X] = ig \int_0^B d\beta' \chi(-i\beta'),$ for $B = f^B - 1(\chi, r^B),$ where χ is the complete Hamiltonian of the system, Eq. (2.1) is transformed to:

$$e_{AB}(v) = -i \int dt \exp(iEt) \langle [A(t), r^B(0)] \rangle. \quad (2.2)$$

$$[X, Y] = XY - YX.$$

In the second quantization and considering that the electron - photon interaction is the principal mechanism of scattering the following holds:

$$\begin{aligned} \chi &= \chi_e + \chi_r + \lambda V; \quad \chi_e = \sum_a \epsilon_a a^\dagger a; \quad \chi_r = \sum_q \mu_q b_q^\dagger b_q; \\ \lambda V &= \sum_{q, aa'} L_q^{aa'} a_q^\dagger a_{a'} (b_q - b_{-q}^\dagger); \quad L_q^{aa'} = C_q(a | \exp(iqr) | a'). \quad \text{For free} \end{aligned}$$

Card 2/13

23103
8/181/83/003/005/008/042
B101/B214

Derivation of a kinetic ...

electrons $L_{\vec{q}}^{aa'} = C_q(n|\exp(i\vec{q}_x \cdot \vec{x})|n')$ $\prod_{j=y,z} \delta(\vec{k}_j - \vec{k}_{j'} + \vec{q}_j)$. The current operators are $\Lambda = \sum_{aa'} \Lambda_{aa'} a_{a'}^+ a_{a'}$. From this follows:

$$\left. \begin{aligned} \sigma_{AB}(E) &= \sum_{aa'} A_{aa'} \tau_{aa'}(E), \\ \tau_{aa'}(E) &= \sum_{q_1 q_2} r_{q_1 q_2}^a G_{q_1 q_2 q_1 q_2}(E). \end{aligned} \right\} \quad (2.3),$$

where

$$G_{q_1 q_2 q_1 q_2}(E) = \langle a_{q_1}^+ a_{q_1} | a_{q_2}^+ a_{q_2} \rangle = \frac{-i}{2\pi} \int d\omega e^{i\omega t} g(t) \langle [a_{q_1}^+(t) a_{q_2}(t), a_{q_2}^+ a_{q_1}] \rangle \quad (1);$$

$g(t) = \begin{cases} 0 & t < 0 \\ 1 & t > 0 \end{cases}$ is the spectral representation of the lagging Green's temperature function. By the method of such equations the equations for $\sigma_{a_1 a_2}^B(E)$ are derived in λ^2 approximation. The following is written down:

Card 3/13

Derivation of a kinetic ...

23103
8/18/61/003/005/008/042
B101/B214

$$(E - \epsilon_{\alpha_1}) G_{\alpha_1 \alpha_2}^{(U)} (E) = f_{\alpha_1} b_{q_1}^+ b_{-q_1}^- + \sqrt{m} b_{q_1}^+ b_{-q_1}^- + (2.4),$$

$$+ \sum_{j=1,3} \sum [L_j^m G_{\alpha_1 \alpha_2}^{(U)} - L_j^m G_{\alpha_2 \alpha_1}^{(U)}].$$

where in the λ^2 approximation

$$v_{\alpha_1 \alpha_2} = b_{q_1}^+ (a_{\alpha_1}^+ a_{\alpha_2}^-) - b_{-q_1}^- (a_{\alpha_1}^+ a_{\alpha_2}^-) \approx (B). \\ \approx f_{\alpha_1} b_{q_1}^+ b_{-q_1}^- + \sqrt{m} b_{q_1}^+ b_{-q_1}^-$$

Here, $f_{\alpha_1 \alpha_2} = f_{\alpha_1} - f_{\alpha_2}$ (f_α = equilibrium distribution of the electrons).
The following holds in Eq. (2.4):

$$G_{\alpha_1 \alpha_2}^{(U)} = \langle a_{\alpha_1}^+ a_{\alpha_2}^- b_q^+ | b_{-q}^- a_{\alpha_1}^+ a_{\alpha_2}^- \rangle. \quad (2.5),$$

where $b_q^{(j=1,3)} = b_q$; $b_q^{(j=2)} = b_{-q}^+$. For nondegenerate electron gas and
Card 4/13

Derivation of a kinetic ...

8/18/61/003/005/008/042
B101/B214

Boltzmann-distribution for a large Gibb's ensemble goes over into canonical ensemble in which the following holds: $\langle a_{a_1}^+ \dots a_{a_s}^+ a_{a_1'} \dots a_{a_r}' | Q \rangle = 0$ for $s > 1, r > 1$ (2.6). From this condition is obtained:

$$(E - \Omega_{\omega_i}) C_{a_1 \dots a_s a_1' \dots a_r'}^{(U)} = C_{a_1 \dots a_s a_1' \dots a_r'}^{(U)} + \\ + \sum_{a_q} [L_{a_q}^{(a)} \langle a_{a_1}^+ a_{a_q} b_{a_q}^{(U)} (b_{a_q} + b_{-a_q}^+) | a_{a_1}^+ a_{a_q} \rangle_s - \\ - L_{a_q}^{(a)} \langle a_{a_1}^+ a_{a_q} (b_{a_q} + b_{-a_q}^+) b_{a_q}^{(U)} | a_{a_1}^+ a_{a_q} \rangle_s], \quad (2.7),$$

where

$$\Omega_{\omega_i}^{(a)} = \epsilon_a - \epsilon_{a_q} - (-1)^q \omega_i, \\ C_{a_1 \dots a_s a_1' \dots a_r'}^{(U)} = \delta_{a_1 a_1'} \langle a_{a_1}^+ a_{a_q} b_{a_q}^{(U)} \rangle - \delta_{a_1 a_1'} \langle a_{a_1}^+ a_{a_q} b_{a_q}^{(U)} \rangle. \quad (c).$$

The following is assumed for the linear electron-photon interaction:
Card 5/13

23103

Derivation of a kinetic ...

S/181/61/003/005/008/042
B101/B214

$$\begin{aligned} & \langle a_1^+ a_2^- b_q^{(U)} (b_{q'} + b_{-q'}^+) | a_1^+ a_2^- \rangle_s \approx \\ & \approx G_{a_1^+ a_2^-}(E) \langle b_q^{(U)} (b_{q'} + b_{-q'}^+) \rangle \approx G_{a_1^+ a_2^-}(E) N_q^{(U)} \\ & N_q^{(U=1, 2)} = N_q = (e^{h\omega_q} - 1)^{-1}, \quad N_q^{(U=\eta)} = 1 + N_q. \end{aligned} \quad (2.8).$$

Introduction of (2.8) in (2.7), calculation of $G_{a_1^+ a_2^- a_2^+ a_1^-}$ and substitution in Eq. (2.4) taking into account (2.3) leads to:

$$(E - \epsilon_{a_1 a_2}) \varphi_{a_1 a_2}^S(E) + L_{a_1 a_2}^S(E) = \Pi_{a_1 a_2}^S(E), \quad (2.9),$$

where

$$\begin{aligned} L_{a_1 a_2}^S(E) = & \sum_{j=1, 2} \sum_q \sum_{\omega} (\varphi_{a_1}^S L_q^{\omega} L_{-q}^{\omega} [N_q^{(U+1)} (E - \Omega_{qj}^{(U)})^{-1} + \\ & + N_q^{(U)} (E - \Omega_{qj}^{(U)})^{-1}] - \varphi_{a_2}^S L_q^{\omega} L_{-q}^{\omega} N_q^{(U)} (E - \Omega_{qj}^{(U)})^{-1} - \\ & - \varphi_{a_1}^S L_q^{\omega} L_{-q}^{\omega} N_q^{(U+1)} (E - \Omega_{qj}^{(U)})^{-1}), \end{aligned} \quad (2.10),$$

Card 6/13

CONT

Derivation of a kinetic ...

S/181/61/003/005/008/042
B101/B214

$$\Pi_{\alpha\alpha}^B(E) = r_{\alpha\alpha}^B \tau_{\alpha\alpha} - \sum_{\alpha'} \sum_{\alpha''} (r_{\alpha\alpha'}^B L_{\alpha'}^{\alpha''} L_{\alpha''}^{\alpha}) [f_q^{(0)}(\alpha', \alpha'') (E - \Omega_{\alpha''}^{(0)})^{-1} + \\ + f_q^{(0)}(\alpha, \alpha'') (E - \Omega_{\alpha''}^{(0)})^{-1}] - r_{\alpha\alpha'}^B L_{\alpha'}^{\alpha''} L_{\alpha''}^{\alpha} f_q^{(0)}(\alpha', \alpha) (E - \Omega_{\alpha''}^{(0)})^{-1} - \\ - r_{\alpha\alpha'}^B L_{\alpha'}^{\alpha''} L_{\alpha''}^{\alpha} f_q^{(0)}(\alpha, \alpha'') (E - \Omega_{\alpha''}^{(0)})^{-1}, \quad (2.11).$$

$$f_q^{(0)}(\alpha, \alpha'') = \frac{\exp(\beta \Omega_{\alpha''}^{(0)}) - 1}{\Omega_{\alpha''}^{(0)}} / N_q^{(0)},$$

$$f_q = \exp[\beta(\mu - \epsilon_q)],$$

Here, $C_{\alpha_1 \alpha_2}^{(j)}$ has been expanded in power series of λ . In strong magnetic field the inhomogeneity of $\Pi_{\alpha_1 \alpha_2}^B(E)$ of Eq. (2.9) contains besides $\Pi_{\alpha_1 \alpha_2}^{(0)} \sim \lambda^0$ also terms $\lambda^2 \Pi_{\alpha_1 \alpha_2}^{(2)B}(E)$ since for $\omega - \omega_{\alpha_2 \alpha_1} \neq 0$ one has: $\text{Re} \sigma_{AB}^{(0)} \sim \lambda^2$. For $\omega_{\alpha_1 \alpha_2}^{(0)B} \sim \lambda^0$ one obtains:

$$(E - \omega_{\alpha_1 \alpha_2}) \tau_{\alpha_1 \alpha_2}^B + f_{\alpha_1 \alpha_2} \tau_{\alpha_1 \alpha_2}^B = \Pi_{\alpha_1 \alpha_2}^B \quad (2.12).$$

Card 7/13

25103

Derivation of a kinetic ...

8/181/61/003/005/C08/042
B101/B214

where $\tilde{\Pi}^B_{[a_1 a_2]} = \tilde{\Pi}^B_{[a_1 a_2]} + R^B_{[a_1 a_2]}$. 2) For free electron gas with effective mass m the zeroeth approximations for $\psi_{n,n+1,n+2}^{(0)}$, $\psi_{n,n+1}^{(0)}$ and $\psi_{n,n+2}^{(0)}$ are introduced in Eq. (2.9) and equations for $\psi_n(k, E) = \psi_{n,k,n+1,k}(E)$ are obtained. For the case $\psi_n(k_z^2, E) = \psi_n(k_z, E)$ the following holds:

$$[(\omega - \epsilon) - Q_n(k_n, \omega)] \Phi_n(k_n) + \sum_{n'=-1,0} \int \frac{dq_n}{(2\pi)^3} Z_{nn'}(k_n, q_n, \omega) \Phi_{n'}(k_n + q_n) = (3.1)$$

$$= i \tilde{\Pi}_n(k_n) - \Pi_n(k_n),$$

where

$$Q_n(k_n, \omega) = \sum_{j=1,2} \sum_{n'=0} \int \frac{(dq_n)}{(2\pi)^3} [|L_q^{n,j}|^2 N_q^{j,0} (\omega - \omega_{n+1,j} + (-1)^j \omega_j) + \\ + N_q^{j+1} |L_q^{n+1,j}|^2 (\omega - \omega_{n+1,j} + (-1)^j \omega_j)], \quad (3.2).$$

$$Z_{nn'}(k_n, q_n, \omega) = \sum_{j=1,2} \int \int \frac{dq_n dq_{n'}}{(2\pi)^6} L_q^{n,j} L_{q'}^{n+1,j} X$$

Card 8/13

Derivation of a kinetic ...

S/181/61/003/005/CC8/042
B101/3214

$$\times [N_q^{U+1}B(u - u_{s+1,s} + (-1)^s u_q) + N_q^UB(u - u_{s+1,s} + (-1)^s u_q)],$$

$$\begin{aligned} \Pi_s''(k_s) = & \Pi_{s,s+1,s}'' - \sum_{j=1,2} \sum_{s'=s}^{\infty} \int \frac{(d\mathbf{q})}{(2\pi)^3} [L_q^{ss'} L_{-q}^{s+1,s'+1} \varphi_{s',s'-1}^{(0)} X \\ & \times [N_q^{U+1}B(u - u_{s',s'-1} + (-1)^s u_q) + N_q^UB(u - u_{s',s'-1} + (-1)^s u_q)] - \\ & - \varphi_{s,s-1}^{(0)} L_q^{s+1,s} L_{-q}^{s,s-1} N_q^{U+1}B(u - u_{s,s} + (-1)^s u_q) - \\ & - \varphi_{s+1,s+1,s}^{(0)} L_q^{s+1,s} L_{-q}^{s+1,s} N_q^UB(u - u_{s+1,s} + (-1)^s u_q)], \end{aligned}$$

$$n' = (n', k' = k + q), \quad \Pi_{s,s+1,s}'' = \lim_{\epsilon \rightarrow 0} \Pi_{s,s+1,s}(E),$$

Here the conservation laws have been taken into consideration: $k' = k + \vec{q}$; $\mathbf{k} = (k_x, k_y)$; $\vec{q} = (\vec{q}_x, \vec{q}_y)$. In the quantum limit for $\beta\omega_0 \gg 1$ Eqs. (3.1) can be solved by the method of successive approximation. For the quantum limit $\beta\omega_0 \gg 1$ (3.1) takes the form:

$$\begin{aligned} & [(u - u_s) - Q_s(k_s, u)] \Phi_s(k_s) + \quad (3.3) \\ & + \int \frac{dq_s}{2\pi} Z_s(k_s, q_s, u) \Phi_s(k_s + q_s) = -\Pi_s''(k_s) + i\Pi_s''(k_s). \end{aligned}$$

Card 9/13

23103

S/181/61/003/005/008/042
B101/B214

Derivation of a kinetic ...

where $Q_0 = Q_{n=0}$; $Z_0 = Z_{n=0, n' \neq 0}$; $\bar{\Gamma}_0^n = \bar{\Gamma}_{n=0}^n$. Hence for the electric conductivity of free electrons one obtains: $\text{Re}\sigma_{xx}(\omega) = e^2 \text{Im}[g(E) - g(-E)]$; $\text{Im}\sigma_{xy}^{(a)}(\omega) = e^2 \text{Im}[g(E) + g(-E)]$; $\text{Re}\sigma_{xy}^{(a)}(\omega) = e^2 \text{Re}[g(E) + g(-E)]$;

$\text{Im}\sigma_{xx}(\omega) = -e^2 \text{Re}[g(E) - g(-E)]$; (3.4), where $g(E) = \sum_{n=0}^{\infty} \int_{-\infty}^{\infty} (dk_z/2\pi) \sqrt{n+1} n k_z \Gamma_{z,z}(E) (\omega_0/q_0)$ (3.5); and for $\beta\omega_0 \gg 1$: $g(E) \approx g_0(E)$

$= \sum_{-\infty}^{\infty} (dk_z/2\pi T^2 q_0 \omega_0^{-1}) \Gamma_0^{(0)}(k_z) q_0(k_z, \omega) + (\omega - \omega_0) \Gamma_0^{(0)}(k_z) / (\omega - \omega_0)^2$. For the case of elastic scattering at acoustic phonons $sq_0 \ll kT$, $\bar{\Gamma}_0 = 0$ and for $g_0(E)$ one obtains:

$\text{Im}g_0(E) = + \int_{-\infty}^{\infty} (dk_z/2\pi T^2 q_0 \omega_0^{-1}) [\Gamma_0^{(0)}(k_z) q_0(k_z, \omega) + (\omega - \omega_0) \Gamma_0^{(0)}(k_z)] / (\omega - \omega_0)^2$
 $+ [q_0(k_z, \omega)]^2]^{-1}$; $\text{Re}g_0(E) = \int_{-\infty}^{\infty} (dk_z/2\pi T^2 q_0 \omega_0^{-1}) (\omega - \omega_0) \Gamma_0^{(0)}(k_z) / (\omega - \omega_0)^2$
 $+ [q_0(k_z, \omega)]^2]^{-1}$ (3.6). By means of Eqs. (3.4)-(3.6), σ_{xx} , σ_{xy} are

Card10/13

23103
S/181/61/003/CC5/CC8/C42
B101/2214

Derivation of a kinetic ...

calculated in the case investigated. A) When for real $k_z \leq m/3$ the following holds: $(\omega - \omega_0)^2 \gg (q_0(k_z, \omega))^2$, one obtains the iteration equations of Ref. 1. B) For $(\omega - \omega_0)^2 \ll (q_0(k_z, \omega))^2 \sim \lambda^4$ in the neighborhood of resonance $(\omega - \omega_0)^2 \sim \lambda^4$ is neglected and the lines of cyclotron resonance for each k_z are a superposition of the quasi Lorenz lines for whose width the following holds: $Q_0(k_z, \omega) = \sum_{l=0,1} W_l(k_z, \omega)$, where

$$W_l(k_z, \omega) = \frac{\pi C_0}{\rho M e^3} \sum_{n=-\infty}^{\infty} \sum_{\pm} \int \frac{(dq)}{(2\pi)^3} |L_n^{\omega}|^2 \delta(\omega + \omega_0(n+l-1) - (-1)^l \Delta \pm sq), \quad (5).$$

C_0 is the constant of the deformation potential, ρ the density of the substance, and $\Delta = (k_z^2/2m) - (k_z + q_z)^2/2m$. In the quantum limit investigated the following holds:

Card 11/13

23103

S/181/61/003/005/008/042
B101/B214

Derivation of a kinetic ...

$$Q_0(k_z, \omega) \propto \frac{q_p^2 m C_0^2}{\pi^2 k_z^3 \sqrt{k_z^2 + 2q_p m \omega}} \begin{cases} \text{spn } 0 < \omega < \omega_0, \\ \text{spn } \omega \approx \omega_0, \end{cases} \quad (3.7).$$

r. e. $Q_0 \sim \frac{1}{k_z^3} (k_z^2 + 2q_p m \omega)^{-1/2}$.

It follows from Eqs. (3.6)-(3.7) that for $\omega = 0$ the logarithmic divergence $\int dk_z/k_z$ of the scattering is removed even without taking into consideration the small inelasticity. If $sq_0 \ll kT$ $Q_0(k_z) \sim |k_z|^{-1}$ is neglected the Born approximation is not valid even for $k_z \rightarrow 0$, and σ_{xx} must be calculated from Eqs. (3.4)-(3.7) taking into account the small inelasticity of $sq_0 \ll kT$. N. N. Bogolyubov and S. V. Tyaplikov are mentioned. There are 15 references: 11 Soviet-bloc and 4 non-Soviet-bloc. The 4 references to English-language publications read as follows: P. Martin, J. Schwinger, Phys. Rev. 115, 1342, 1959; R. Kubo, Phys. Soc. Jap., 12, 570, 1957; E. Adams, T. Holstein, J. Phys. Chem. Sol., 10, 254, 1959; A. Kahn, Phys. Rev., 119, 1960.

Card 12/13

Derivation of a kinetic ...

23103
S/181/61/UU2/005/008/042
B101/B214

ASSOCIATION: Institut poluprovodnikov AN SSSR Leningrad (Institute of Semiconductors AS USSR, Leningrad)

SUBMITTED: August 22, 1960 (initially), February 4, 1961 (after revision)

Card 13/13

KLINGER, M.I.

Author's correction to his article "Theory of linear irreversible processes in a high magnetic field" ("Fizika tverdogo tela," vol. 3, p. 1342, 1961). Fiz. tver. tela 3 no.8:2577-2508 '61.

(MIRA 14:6)

(Irreversible processes)
(Magnetic fields)

24.7100 (1144, 1385, 1559)

30062
S/048/61/025/011/006/031
B108/B138AUTHOR: Klinger, M. I.

TITLE: Theory of the kinetic phenomena in semiconductors of the NiO type

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Seriya fizicheskaya, v. 25, no. 11, 1961, 1342-1346

TEXT: Electron transfer in semiconductors of the NiO type has a number of peculiar features at high temperatures the most important of which is particularly low mobility $U(T)$. In order to elucidate these, the author develops a general theory of transfer with low mobility. The system considered is a small polaron as a carrier at a lattice site in a transition metal which interacts weakly with the remaining, stationary ions.

A. F. Ioffe (Canad. Phys., 34, 1393 (1956)) had originated the idea that carriers jump from one lattice site to the other. This also holds here if the polaron band width $\Delta_{\text{pol}} < \hbar\Gamma(T)$, where $\Gamma(T)$ is the mean width of the unperturbed polaron level, i.e., small polarons and continuous phonon spectrum. The theory is built up on the basis of previous papers

Card 1/3

30062

S/048/61/025/011/006/031

B108/B138

Theory of the kinetic phenomena in...

(Klinger M. I., Fizika tverdogo tela, 1, 1225 (1959); 2, 309 (1960)) in which the author defined the kinetic coefficients σ_{AB} . In this paper he considers only steady processes for which $\sigma_{AB}^{(s)} = \frac{1}{2} (\tilde{\sigma}_{AB} + \tilde{\sigma}_{BA})$, where

$$\tilde{\sigma}_{AB} = \frac{BN}{2} \int_{-\infty}^{\infty} dt e^{-i\omega t} \text{Re} \langle BA(t) \rangle. \quad (2)$$

B and A denote the fluxes of charge, energy, etc. The Hamiltonian of the system under review is $H = H_0 + \lambda_0 H'$ with

$$H_0 |sn^{(s)}\rangle = s(n^{(s)})^2 |sn^{(s)}\rangle; \langle sn^{(s)}|H'|s'n'^{(s')} \rangle = \langle sn^{(s)}|V_r|s'n'^{(s')} \rangle; \quad (3)$$

$$\langle sn^{(s)}|H'|sn^{(s)}\rangle = 0; \lambda_0 V_r = \sum_{r' \neq r} V_r(x - ar);$$

$\langle \dots \rangle = \text{Sp exp}(\beta P - \beta H)(\dots)$; N_0 is the carrier concentration; $\beta = 1/kT$. $n^{(s)}$ is the number of phonons, which may vary in an $s \rightarrow s'$ transition (s, s' are lattice nodes). $\langle sn^{(s)}|s'n^{(s')} \rangle = \delta_{ss'} \delta_{nn'}$. The term $\lambda_0 H'$ is

Card 2/3

Card 3/3

43936

0/030/62/002/008/003/003
I046/1246

24.2700

AUTHOR:

Klinger, M.I.

TITLE:

The theory of transport phenomena in low-mobility semiconductors
(Derivation of formulas for fundamental kinetic coefficients)

PERIODICAL:

physica status solidi; v. 2, no. 8, 1962, 1062-1087

TEXT: This work is a sequel to M.I.Klinger's previous papers (Ref.1: Izv. Akad. Nauk SSSR, physical series, no. 11, 1942, 1961; Ref.2: DAN SSSR, 142, 1065, 1962) where the general theory of transport phenomena in low-mobility semiconductors is developed without reference to any concrete model. It is shown that transport involves multi-phonon transitions of polaron-type carriers between neighboring lattice cells. The general methods of Ref. 2 are used in determining the electric conductivity and thermoelectric and thermomagnetic coefficients. The energy transport due to polarons is found to be negligible. It is also shown that drift involves independent transitions whereas the Hall effect is determined by phase-correlated transitions. All the results are obtained for crystals where the concentration of lattice cells is much higher than

Card 1/2

The theory of transport phenomena...

the concentration of carrier and impurity.

ASSOCIATION:

Institut poluprovodnikov Akademii Nauk SSSR Leningrad (The Institute
of Semiconductors, Academy of Sciences USSR, Leningrad)

SUBMITTED:

May 29, 1962

Card 2/2

S/181/62/004/011/009/049
B102/B104

AUTHOR: Klinger, M. I.

TITLE: Theory of transfer effects in semiconductors with low mobility
(the characteristics of the electric spectrum of the system)

PERIODICAL: Fizika tverdogo tela, v. 4, no. 11, 1962, 3075 - 3085

TEXT: A theoretical investigation of the energy spectrum of ordered and disordered semiconductors with low mobility is described, $u \ll \frac{e}{m}$,
 $a^2 / h \text{ cm}^2/\text{v.sec}$ where $|e|a^2 / h \approx 1 \text{ cm}^2/\text{v.sec}$ at $a \approx 3 \cdot 10^{-8} \text{ cm}$, a is the lattice constant or in general the distance of adjacent ions. Starting from a system of interacting phonons and stripped carriers (electrons, holes, or Frenkel' excitons) located in the static field of a semiconductor, the general properties are investigated for an unperturbed system and for a system perturbed in its $|\Psi_n\rangle$ base. Any anharmonic phonon characteristic is neglected. A representation of the orthonormalized $|\Psi_n\rangle$ base is obtained for localized carrier functions and for the spectrum. The complete perturbation matrix differs from that given in Phil. Mag. 3, 1361, 1958; J. Phys. Card 1/4

Theory of transfer effects

S/181/62/004/011/009/049
B102/B104

Chem. Sol. 5, 34, 1958; Ann. of Phys. 8, 343, 1952. The singularities are studied. The general case for $n=n'$ yields

$$\langle sn | \mathcal{H}' | s'n' = n \rangle = \left\{ (s | \delta V | s') + \sum_j [(X_{ij}^s)^* N_{ij} - X_{ij}^s (1 + N_{ij})] V_{ij}^s + \right. \\ \left. + \sum_j [X_{ij}^s N_{ij} - X_{ij}^s (1 + N_{ij})] (V_{ij}^{s'})^* \right\}_{s \neq s'} \exp(-\Phi_{ss'}(n)); \quad (2.30).$$

For $H = 0$ gives

$$\langle sn | \mathcal{H}' | s'n' = n \rangle = \langle s'n' | \mathcal{H}' | sn \rangle, \text{ and}$$

$$(2.31) \quad \begin{aligned} \langle sn | \mathcal{H}' | s'n' = n \rangle &= \delta V^\omega (s^n | s'^n), \\ \delta V^\omega &= (s | \delta V | s' + \sum_j [V_{ij}^s (X_{ij}^s)^* + (V_{ij}^{s'})^* X_{ij}^s]). \end{aligned}$$

$V_{ij}^s(x)$ is the interaction coefficient of a dressed carrier with phonons with the quasi-momentum \vec{k} of the j -th branch and with a frequency ω_{ij}^s , N_{ij}^s is the number of phonons, δV the potential fluctuation,

$$V_{ij}^s \equiv V_{ij}; V_{ij}^\omega \equiv (s | V_{ij}(x) e^{i\omega x} | s), \quad X_{ij}^s \equiv -V_{ij} \cdot \omega_{ij}^{-1} \sim N_{ij}^{-1}.$$

The investigation of an unperturbed system and of a perturbation in the (kn) base follows. The interpolation of the system characteristics in the

Card 2/4

Theory of transfer effects ...

S/181/62/004/011/009/049
B102/B104

$|kn\rangle$ and $|k'n\rangle$ bases is studied. The orthonormalized $|kn\rangle$ base of non-localized carrier functions is represented, in taking the translation symmetry of the crystal into account. The expressions obtained for the carrier energy in the bands differ by a pre-exponential factor from those given in the papers mentioned above. The following expressions are derived:

$$\delta_n(k) = \sum_{s' \neq s} \delta V(s-s') \cos k_n(s-s') \exp[-\Theta_\omega(s)], \quad (3.6)$$

$$\Theta_\omega(s) = \sum_{ij} \frac{1}{2} |X_{ij}^\omega|^2 (1 + 2N_{ij}). \quad (3.7)$$

$$k_n = k - \sum_{ij} f N_{ij}, \quad \text{carrier quasi-momentum}$$

$$(kn|\mathcal{H}|k'n) = \delta_{kk'} N_0^{-1} \sum_{n'n} e^{-ik(n'-n)} (kn|\mathcal{H}'|s'n), \quad (3.16)$$

$$(kn|\mathcal{H}'|k'n) \sim \delta_{kk'} N_0^{-1/2} |f|_{>1} n p n, n \neq n'. \quad (3.17)$$

$$\text{Card } 3/4 \quad J_{kn,k'n}^{(0)} \sim (kn|\mathcal{H}'|f_1(\mathcal{H}_0) \cdot \dots \cdot f_{l-1}(\mathcal{H}_0) \mathcal{H}'|k'n) \sim \begin{cases} \delta_{kk'} \delta_{nn'} \\ \delta_{kk'} (1 - \delta_{nn'}) N^{-1/2} \end{cases} \quad (3.18)$$

Theory of transfer effects . . .

S/181/62/004/011/009/049
B102/B104

$$\begin{aligned}
 \langle sn | \mathcal{H}' | s'n' \rangle &= \left\{ \delta V^{ss'}(n^{(0)} | n'^{(0)}) + \sum_{ij} [V_{ij}^{ss'}(n^{(0)} | C_{ij,s} | n'^{(0)}) + \right. \\
 &\quad \left. + (V_{ij}^{ss'})^*(n^{(0)} | C_{ij,s}^* | n'^{(0)}) \right\}_{s,s'} \exp[i(x_s s - x_{s'} s')] = \\
 &= \left\{ (s | \delta V | s') (n^{(0)} | n'^{(0)}) + \sum_{ij} [V_{ij}^{ss'} \prod_j (0 | (b_{r,s})^{r,s} (T_{rr}^{s'})^* b_{r'} | 0)]_s \times \right. \\
 &\quad \left. \times T_{r,s'}^* (b_{r,s'}^*)^{r,s'} | 0 \rangle - \langle V_{ij}^{ss'} \prod_j (0 | (b_{r,s})^{r,s} (T_{rr}^{s'})^* b_{r'} | T_{r,s'}^* (b_{r,s'}^*)^{r,s'} | 0) \rangle \right\}_{s,s'} \times \\
 &\quad \times \exp[i(x_s s - x_{s'} s')]. \tag{2.23}
 \end{aligned}$$

Finally the relation of $\langle \hat{s}n |$ and $\langle \hat{k}n |$ functions and expressions for complex term shifts in both bases are examined.

ASSOCIATION: Institut poluprovodnikov AN SSSR, Leningrad (Institute of Semiconductors AS USSR, Leningrad)

SUBMITTED: March 12, 1962 (initially)
June 2, 1962 (after revision)

Card 4/4

8/101/62/004/011/010/049
B102/B104

AUTHOR:

Klinger, M. I.

TITLE:

Theory of transfer effects in semiconductors with low mobility
(general approach in the theory of transfer processes and
criteria of this theory)

PERIODICAL: Fizika tverdogo tela, v. 4, no. 11, 1962, 3086 - 3103

TEXT: The theory of transfer effects is extended, and its limits determined, in continuation of earlier investigations (PTT, 4, 11, 3075, DAN SSSR, 140, no. 5, 1065, 1962; Izv. AN SSSR, ser. fiz. No. 11, 1342, 1961). The object was to arrive at general basic relations enabling any kinetic coefficient to be calculated and also determine the criteria for applicability of this theory. The basic equations for calculating the kinetic coefficients are derived, and for the zeroth approximation the following expression is obtained:

$$\sigma_{xx}^{(0)} = \beta \int dt \frac{1}{2} \sum \epsilon^{\mu\nu\rho\sigma} \text{Re} (N_\omega(\epsilon_i' | M | \epsilon_i) + M_\omega(\epsilon_i' | N | \epsilon_i)), \quad (2,36)$$

where
Card 1/3

Theory of transfer effects in ...

S/181/62/004/011/010/049
B102/B104

$$|\epsilon_i\rangle \equiv \exp[-(E_n f - \Gamma_n f)/\epsilon] |\epsilon\rangle$$

$$\Gamma_e = \sum_{n'} |(\epsilon_n | \mathcal{K}' | \epsilon_{n'})|^2 \Gamma_n(n') [(\epsilon_n - \epsilon_{n'})^2 + (\Gamma_n(n'))^2]^{-1} \quad (2.37).$$

An example is given showing how to obtain the first and second order correction formulas. The first-order correction is of the following type

$$(\epsilon | N | \epsilon') (\epsilon' | \mathcal{K}' | \epsilon'') (\epsilon'' | N | \epsilon). \quad (2.40).$$

The limits of application are investigated for semiconductors with a strong electron-phonon coupling ($\Phi_0 \gg 1$) and with a small radius of the polaron-type carriers. The contribution of each approximation to various special cases is estimated. Two appendices deal with problems arising in these approximations. Conclusions: In order to realize the transfer effects it is necessary to have a high enough temperature, i. e. $T > T_0$, where $T_0 \approx \omega_p$ and ω_p is the characteristic phonon frequency. Another necessary condition is $\Delta_e / \Phi_0 \omega_p \gg 1$ where Δ_e is the electron band width. It follows from the criteria that at $T > T_0$ a new transfer type will occur, caused by transitions of localized carriers between lattices. Uneven effects with respect to the

Card 2/3

Theory of transfer effects in ...

S/181/62/004/011/010/049
B102/B104

magnetic field, such as the Hall effect, are determined by the contribution of phase-correlated transitions. In this respect the transfer shows a quantum-type character.

ASSOCIATION: Institut poluprovodnikov AN SSSR, Leningrad (Institute of Semiconductors AS USSR, Leningrad)

SUBMITTED: March 12, 1962 (initially)
June 2, 1962 (after revision) ✓

Card 3/3

34823

14,7700(1035,1043,1055)

8/020/62/142/005/014/022
B104/B102

AUTHOR: Klinger, M. I.

TITLE: Theory of kinetic effects for low carrier mobility

PERIODICAL: Akademiya nauk SSSR. Doklady, v. 142, no. 5, 1962,
1065 - 1068

TEXT: A. F. Ioffe (Canad. J. Phys., 34, no. 12A, 1393 (1956)) showed the carrier mobility of some semiconductors to be anomalously low. The transport mechanism in these semiconductors was shown to differ largely from the usual one. The author formulates a theory free of limitations and certain difficulties for transport effects in an ideal crystal with small polarons or other small polarizable carriers. Electron-phonon interaction is assumed and an electron (hole) in the field of vibrating ions of a three-dimensional ionic lattice is studied. The total Hamiltonian is given as the sum of the Hamiltonian of the undisturbed system and that of small disturbances. The transport effects are studied in a system with strong electron-phonon coupling. The kinetic

Card 1/3

J/020/62/112/005/014/022
B104/B102

Theory of kinetic...

coefficients are calculated by expressions from preliminary studies (M. I. Klinger, *Fiz. Tverd. tela*, 11, no. 12, 1342 (1960); E. Kubo, *J. Phys. Soc. Japan*, 12, no. 1, 570 (1957)). The condition $\gamma \equiv \Delta_c^{\text{mean}} / \omega_e \ll 1$, (i.e., γ and Δ_c^{mean} are the small parameters of the theory) is a criterion of the applicability of the perturbation theory to the calculation of Δ_{AB} [Abstracter's note: Δ_{AB} not defined]. $\Delta_c^{\text{mean}} = \Delta_c$ $\exp(-\phi(T))$, where Δ_c is the carrier band width, ω_e the characteristic frequency phonon. $\gamma \ll 1$ means that the migration occurs due to transitions of localized polaron-type wave packets between the cells. This takes place only if the lifetime of a packet in a cell is shorter than its

Card 2/3

Theory of kinetic...

S/020/62/142/005/014/022
B104/B102

leakage time. } is the condition of quasisteadiness of the state of polarization of the lattice around the carriers. This paper was the subject of a report given at the Soveshchaniye po ferro- i antiferromagnetizmu (Conference on Ferro- and Antiferromagnetism) in Leningrad on May 6, 1961. There are 11 references: 4 Soviet and 9 non-Soviet. The four most recent references to English-language publications read as follows. A. Miller, E. Abrahams, Phys. Rev., 120, no. 3, 745 (1960); C. H. Jonker, J. Phys. Chem. Sol., 9, no. 27, 165 (1959); L. van Houten, Phys. Chem. Sol., 17, no. 1/2, 7 (1960); T. Holstein, Ann. of Phys., no. 3, 343 (1959). ✓

ASSOCIATION: Institut poluprovodnikov Akademii nauk SSSR (Institute of Semiconductors of the Academy of Sciences USSR)

PRESENTED: July 24, 1961, by A. A. Lebedev, Academician

SUBMITTED: June 28, 1961

Card 3/3

L 14391-63
AT/IJP(C)

EWT(1)/EWG(k)/BDS/EEC(b)-2 APPTC/ASD/ESD-3 Pg-4

ACCESSION NR: AP3001728

9/0030/63/003/003/0805/0823

66

62

AUTHOR: Klinger, M. I.

TITLE: Hall effect in low mobility semiconductors)

SOURCE: Physica status solidi, v. 3, no. 5, 1963, 803-823

TOPIC TAGS: Hall mobility, electrophonon coupling, transport phenomena, mobility tensor, semiconductor, Hall angle

ABSTRACT: Expressions are derived for the antisymmetric part of the mobility tensor (to a first, linear, order approximation in magnetic field) and for the Hall mobility (Hall angle) for a low mobility semiconductor. The purpose of this and previous works by the author is the developing of a simple variation of the theory of polaron transport of the non-boltzmann type, for which the usual transport theory is not applicable. An estimate is made of the magnitude and dependence of the Hall mobility on temperature and on the electron-phonon coupling constant. Further calculations show that the Hall mobility is usually much larger than the drift (ohmic) mobility.

Card 1/2

L 14391-63

ACCESSION NR: AP3001728

"The author expresses deep thanks to Professors M. A. Krivoglaz, I. M. Lifshits and S. I. Pekar for their discussions about the problem of the classical limit which were a stimulus to its more detailed analysis; the author likewise acknowledges the assistance of Prof. S. I. Pekar with whom he discussed the contribution of excited states which stimulated more detailed analysis of that problem." Orig. art. has: 52 equations.

ASSOCIATION: Institut poluprovodnikov AN SSSR, Leningrad (Semiconductor Institute of the Academy of Sciences of the SSSR)

SUBMITTED: 05Nov62

DATE ACQ: 10Jun63

ENCL: 00

SUB CODE: GE

NO REF Sov: 004

OTHER: 007

Card 2/2

L 12879-63
AT/IJP(C)

BNT(1)/EMO(k)/MDS/XRD(b)-2

ATTC/ASD/ESD-3

P2-4

ACCESSION NR: AP3000513

8/0020/63/150/002/0286/0289

63

AUTHOR: Klinger, M. I.

62

TITLE: Theory of low-temperature transfer in semiconductors with low mobility.

SOURCE: AN SSSR. Doklady, v. 150, no. 2, 1963, 286-289

TOPIC TAGS: low-temperature transfer, semiconductor, non-Boltzmann transfer, polaron

ABSTRACT: The first part of the article discusses the theory of high temperature transfer in semiconductors of low mobility, i.e. some approximate solution of the problem of a non-Boltzmann transfer in an electrophone system with intense reaction, whose parameter PHI is much greater than 1. The transfer coefficients Sigma sub MN are normally computed by a general formula which is typified by equation (1) of enclosure 1. Corrections for ReD sub MN are discussed in detail. The second part of the article examines the basic correlations of this transfer theory in the same semiconductors, but at low temperatures. In this region of T, using a static case for simplicity, for Sigma sub MN sup (s, a) being identically equal to 1/2 (Sigma sub MN + or - Sigma sub NM), equations (2) and (3) of enclosure 2 are valid. Author then discusses equations for describing carriers (polarons) and presents modifications for equations (2) and (3). Orig. art. has: 23 equations.

Cord. 1/A

Inat. of Semi-Conductors

ACCESSION NR: AP4042789

8/0020/64/157/003/0566/0569

AUTHOR: Klinger, M. I.

TITLE: Theory of nonstationary conductivity of semiconductors
with low mobility

SOURCE: AN SSSR. Doklady*, v. 157, no. 3, 1964, 566-569

TOPIC TAGS: semiconductor carrier, carrier mobility, semiconductor conductivity, polaron, crystal lattice

ABSTRACT: The theory developed by the author elsewhere (Phys. tverd. tela, a) v. 4, 3075, 1962; b) 4, 386, 1962; Phys. Stat. Sol. v. 2, 1062, 1962; DAN v. 142, 1065, 1962; Izv. AN SSSR ser. fiz., no. 11, 1342, 1961) is used to calculate the tensor of nonstationary conductivity $\sigma_{\mu\nu}(\omega)$ and mobility $u_{\mu\nu}(\omega)$ for a semiconductor with low mobility, in which a carrier such as a small polaron is described

Card 1/2

ACCESSION NR: AP4042789

essentially by a local-type function. Approximations are given for the ohmic mobility and for the contributions made to it by the "jumps" of the polaron packet and its "diffusion." It is deduced that the mobility has a noticeable maximum as a function of the frequency. The Faraday angle and the mobility are determined for a magnetic field parallel to the OZ axis in a similar fashion for not too small frequencies, for crystals in which three suitable sites can be mutually nearest neighbors. It is shown that usually the Faraday angle decreases with increasing frequency and with increasing temperature. Orig. art. has: 11 formulas. Report presented by A. A. Lebedev.

ASSOCIATION: Institut poluprovodnikov Akademii nauk SSSR (Institute of Semiconductors, Academy of Sciences SSSR)

SUBMITTED: 07May63

ENCL: 00

SUB CODE: 88

NR REF Sov: 005

OTHER: 005

Card 2/2

L 20580-66

ACC NR: AP6002041

SOURCE CODE: GE/0030/65/012/002/0765/0774

AUTHOR: Klinger, M. I.

55
73ORG: Institute of Semiconductors of the Academy of Sciences of the
USSR, LeningradTITLE: Theory of transport phenomenon in low mobility crystals. II.
Relaxation and transport of small-radius polarons as a stochastic
jumping process

SOURCE: Physica status solidi, v. 12, no. 2, 1965, 765-774

TOPIC TAGS: transport phenomenon, semiconductor crystal, crystal
property, crystal theory, polaron, stochastic process, mass energy
relation, physical diffusion, Boltzmann distribution, Markov processABSTRACT: A stochastic theory of high temperature of small polaron
transport was developed by generalizing on Sevill's model (G. Sevill,
Phys. Rev. 129, 597, 1963). Relaxation of local small polarons is
discussed, and a quantitative picture of this non-Boltzmann type of
transport is given. The transport and relation of such carriers can
be considered as a stochastic type of hopping diffusion, but its main
equation is non-Markovian (with "memory"). Orig. art. has: 19 formulae.
[Based on author's abstract]

SUB CODE: 20/ SUBM DATE: 22 Mar 65/ ORIG REF: 002/ OTH REF: 009

[NT]

Card 1/2 OK

KLINGER, M.I.

Characteristics of the transfer and relaxation of small-radius polarons. Dokl. AN SSSR 165 no.3:520-523 N '65.

(MIRA 18:11)

1. Institut poluprovodnikov AN SSSR. Submitted February 16, 1965.

KLINGER, O., ing.

Strengthening a metallic railroad superstructure with a
prestressed tie rod by heating. Rev sailer for 12 no. 6:
322-326 Je '64.

1. Planning Institute for Transport and Telecommunications.

KLINGER, P.

"Mechanization of Harvesting Coarse Fodder. p. 353 (JARNUTEK ES GEPEK
Vol. 1, No. 12, Dec 1954; Budapest, Hungary.)

So: Monthly List of East European Accessions, (EEL), LC, Vol. 4, No. 4,
April 1955, Uncl..

KLINGER, P.

KLINGER, P. Possibilities of mechanizing maize production. p. 218

Vol. 8, no. 5, May 1956

AGRATUKOMANY

AGRICULTURE

Budapest, Hungary

See: East European Accession, Vol. 6, No. 3, March 1957

KLINGER V.G.

Category : USSR/Optics - Physical Optics

K-5

Abs Jour : Ref Zhur - Fizika, No 2, 1957, No 4999

Author : Vulis, L.A., Klinger, V.G.

Title : Concerning the Problem of the Calculation and Simulation of Radiant Heat Exchange.

Orig Pub : Zh. tekhn. fiziki, 1954, 24, No 11, 2070-2078

Abstract : The equations for the radiant heat exchange between gray bodies, separated by a medium that is transparent to rays, are considered. A computation procedure is proposed, based on the direct connection between the intrinsic and resultant radiation. The possibilities of using a light-ray analogue of the radiation heat exchange are evaluated.

Card : 1/1

KLINGER, V.G.

USSR/Optics - Physical Optics,

K-5

Abs Jour : Referat Zhur - Fizika, No 3, 1957, 7785

Author : Klinger, V.G.

Inst :

Title : Experiment with a Light Analogue of Radiant Heat Exchange

Orig. Pub : Zh. tekhn. fiziki, 1954, 24, No 11, 2083-2089

Abstract : A discussion of the problem of simulating by means of light the radiant heat exchange in a system of opaque gray bodies, separated by a transparent medium. Each body in the system has equal values of the temperature and of the absorbing ability (A) over the entire surface. The surfaces have only diffused reflection and radiation. The purpose of the work was to determine the flux of the resultant radiation (Q) on the surface (F), connected with the density of radiation by the formula $a = dQ/dF = AE_{inc} - E$, where E is the density of the intrinsic radiation, and E_{inc} is the density of the flux incident on the

Card 1/3

- 64 -

USSR/Optics - Physical Optics.

K-5

Abs Jour : Referat Zhur - Fizika, No 3, 1957, 7785

given surface. The surfaces of the model were made illuminated so that the distribution of the temperature on the surface is specified by the corresponding distribution of the luminosity in accordance with the Stefan-Boltzmann law. In this case the outer sphere was non-illuminated. The internal spherical illuminator was made of ground glass. The investigated surfaces had non-selective reflection. The flux densities were measured with selenium photocells. The constancy of the voltage on the tubes, and consequently of the luminosity, was insured by means of rheostats. The first series of experiments was carried out at various values of intrinsic radiation. In the second series of experiments investigation was made of the dependence of the resultant flux on the geometric dimensions and on the optical properties of the surfaces of the model. The values of the resulting fluxes, obtained experimentally, are in good agreement with

Card 2/3

- 65 -

USSR/Optics - Physical Optics.

K-5

Abs Jour : Referat Zhur - Fizika, No 3, 1957, 7785

with those calculated by the well-known Christiansen formula.

The discrepancy between the experimental data and the theory amounts to from 2 to 8%.

Card 3/3

- 66 -

KLINGER, V. G.


KLINGER, V. G.--"Investigation of the Radiation Exchange of Energy Using a Light Model." Kazakh State U imeni S. M. Kirov. Physicomathematical Faculty. Alma-Ata, 1955. (Dissertation for the Degree of Candidate of Physicomathematical Sciences).

SO: Knishnaya Letopis' No. 2, 2 July 1955

"APPROVED FOR RELEASE: 09/18/2001

CIA-RDP86-00513R000723130008-7

KITINGER V.G.



APPROVED FOR RELEASE: 09/18/2001

CIA-RDP86-00513R000723130008-7"

USSR/Physiology of Plants - Photosynthesis.

I.

Abs Jour : Ref Zhur - Biol., No 15, 1958, 67788
Author : Darchiyan, Sh.P., Kurmycova, A.Kh., Klinger, V.G.
Inst : Academy of Sciences KazSSR.
Title : A Comparison of the Spectral Luminosity of Live and Torn-Off Plant Leaves.
Orig Pub : Tr. Sektora astrobotan. AN KazSSR, 1957, 5, 174-186.
Abstract : Photographs were taken of the reflection spectra of leaves of the second stratum of lilac, jasmine, and wild mallow; then the leaves were removed from the plants and photographed immediately. Additional photographs were taken after 5, 10, 20, and 40 minutes, one hour, and two hours. Several series of spectrograms of gypsum and barite screens served as a photometric scale. Standard and ultra-violet spectrographs were used with a glass optic.

Card 1/2

USSR/Physiology of Plants - Photosynthesis.

I.

APPROVED FOR RELEASE: 09/18/2001 CIA-RDP86-00513R000723130008-7
Abs Jour : Ref Zhur - Biol., No 15, 1958, 67788

From a comparison of the course of spectral curves for live and torn-off leaves it was found that in the course of the day there were no important variations in the optical characteristics of the leaf, regardless of when it was torn off the plant. The spectral curves were also compared for sunlight and artificial illumination. On the basis of the data acquired the authors consider that by studying the leaves torn off the plant under artificial illumination, it is possible to determine the complete light balance of the plants, to examine the plants in any weather and regardless of their place of growth, to trace the 24-hour course of photosynthesis by using the curves of spectral luminosity of the plants, and to conduct parallel experiments by the spectro-analysis and physiological methods.
-- I.B. Sharovatova.

Card 2/2

24.5200

9/058/61/300/008/020/044
A055/A101

AUTHOR: Dubovik, I. I., Klinger, V. G.

TITLE: The light transfer between mirror and diffusion surfaces

PERIODICAL: Referativnyy zhurnal, Fizika, no. 8, 1951, 176, abstract 83195
(v sb. "Issled. protsessov perenosu. Vopr. teorii otnositel'nosti".
Alta-Ata, 1959, 97-100)

TEXT: On the model of light the author solves the problem of the radiant transfer between two mirror surfaces and between a mirror and diffusion surface.

[Abstracter's note: Complete translation]

✓B

Card 1/1

PAGE 1 THIS INFORMATION 807/379

American Statistical Association
Mathematics Education Project, "Report on World Scientific Events (Study of Trends for Progress). Trends in the Theory of Relativity" (Albuquerque, NM), 1970, pp. 5-12.
American Mathematical Society, "Relativity: 1970," 1,000 copies printed. (Series: The Trinity)

Mathematical Society U.S.A., Rutherford, E.B., Barnes, and S.M. Patterson, Phys. Rev. Letters 19, 1967.

Mathematical Society U.S.A., Rutherford, E.B., Barnes, and S.M. Patterson, Phys. Rev. Letters 19, 1967.

This collection of articles is intended for research physicists and mathematicians. It can also be used by lecturers and students at universities.

This collection of articles contains the results of 19 studies in theoretical physics and the general theory of relativity made from 1965 to 1970 by the staff of the Institute of General Relativity, School of Mathematics and Mathematics, Inc., Salt Lake City (Department of General Physics and Department of the U.S. Army Research Staff University). The articles are arranged in two groups. Group one contains 15 articles concerning the physical aspects of the topics of quantum mechanics, particle theory, field theory, and the theory of the properties of general relativity. Group two contains 14 articles concerning the properties of matter, particle and wave mechanics, or nuclear models and methods. Group three contains 10 articles on the evolution of the properties of generalized relativity. Article one of the collection is an introduction to the mathematical and physical foundations of quantum mechanics and gives a fairly detailed history of the development of quantum mechanics. References, bibliography, and notes are included.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

II. MODELS OF THE HISTORY OF RELATIVITY

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F., and V.L. Smith, "Statistical Methods for Data Analysis," 1973, 2nd ed., Springer-Verlag, Berlin.

Chernoff, F.

87758

24.5200 (1191, 1498, 1537) S/096/61/000/002/011/014
E194/E155

AUTHORS: Vulis, L.A., Doctor of Technical Sciences,
Gurvich, A.M., Doctor of Technical Sciences, and
Klinger, V.G., Candidate of Technical Sciences

TITLE: Optical Modelling of Radiant Heat Exchange in Furnaces

PERIODICAL: Teploenergetika, 1961, No.2, pp. 67-71

TEXT: The general theory of similarity requires that an optical model should fulfil the three conditions: geometrical similarity; identity of optical properties of surfaces and media; and similarity in the distribution of radiation sources. A special feature of the optical modelling method developed in the Kazakh University is that it avoids fulfilling the third condition by determining on the model a system of optical-geometrical parameters. When these are known it is possible to calculate the distribution of radiant fluxes with an arbitrary distribution of sources in the system in which only the first two conditions need be observed. Thus the technique of optical modelling is greatly simplified. The object of the present article is to direct attention to this method which is still not sufficiently widely used. Accordingly,

Card 1/6

W

87758
8/096/61/000/002/011/014
E194/E155

Optical Modelling of Radiant Heat Exchange in Furnaces
the essentials of the method are described and practical results are given. In optical modelling of radiant heat exchange the radiant fluxes are so low that the temperature factor does not enter into the experiment. The method is nevertheless applicable to studies of furnaces where heat fluxes and temperatures are high, because the equations of radiant heat exchange are the same whatever the energy or spectral composition of the radial fluxes. The temperature distribution is determined in the model by the self-radiation distribution both in the volume and on the walls. From this the temperature distribution is calculated on the basis of the Stefan-Boltzmann law if an integral radiation is modelled; or by Wien's formula if the nature of radiation is being studied from separate spectral bands. The present article considers only integral radiation and assumes that the radiating walls and media have the properties of grey diffuse radiating and absorbing bodies. In modelling, the object is sub-divided into a number of surface and volume isothermal zones. The optical properties of the surface zones are characterised by the mean absorption capacity and those

Card 2/6

44

A7758

S/096/61/000/002/011/014
E194/E155

Optical Modelling of Radiant Heat Exchange in Furnaces
of the volumetric zones by the attenuation factor of the medium.
Modelling consists in constructing a geometrically similar system
of surfaces having optical properties identical with the original
and a similar distribution of isothermal zones. The attenuating
properties required of the medium are discussed. The method is
based on the principle of additivity of radiant fluxes which makes
it possible to determine the optical-geometrical parameters of the
model. If only one surface zone in the model is radiating, the
incident flux on different elements of surface can be measured.
These incident fluxes will be both those received by direct but
attenuated radiation and those reflected from other surfaces. By
successively making one zone luminous after another and measuring
the resultant incident fluxes, dimensionless absorption factors may
be determined for various elements considered. Then the results
are summated to determine the flux density incident on any zone
from all the other zones. Simple formulae are derived and it is
shown that by tests on a single model and simple calculations it is
possible to solve a range of problems. The study of radiant heat

Card 3/6

44

87758
8/096/61/000/002/011/014
E194/E155

Optical Modelling of Radiant Heat Exchange in Furnaces

exchange in the chamber of a stoker-fired furnace by optical modelling is then considered. A transparent plastic model in the shape of a cube of side 15 cm inside contained thin metal walls painted matt black. In one of the walls there were 64 holes which were used to measure the incident radiation. Various difficulties that arise in making the measurements are described. Experience has shown that they can be largely overcome if a thin layer of translucent celluloid with a matt surface is placed between the inner wall of the model and the outer. As incident radiation may be at any angle, the sensitivity of the pick-up should not depend on the angle of incidence. This condition is largely satisfied by a germanium photodiode operating as a valve. This photodiode has maximum sensitivity in the infrared where the absorbing capacity of ordinary water is fairly great. Accordingly water may be used as the attenuating medium in the model. The problem of modelling self-radiation of the medium filling the volume is overcome by having a single source of radiation, moving it from place to place and summatting the results. The particular model described was divided

Card 4/6

LH

67758

S/096/61/000/002/011/014
E194/E155

Optical Modelling of Radiant Heat Exchange in Furnaces
into 6⁴ zones. The radiating element was a cube of transparent plastic, corresponding accurately to the size of the zone. The cube contained a small lamp; it was filled with water and the outside was covered with translucent celluloid. To check the experimental procedure a model was used to study the radiant output of a cylindrical source for which a method of calculation exists. The calculated and experimental results agreed within 5% and the accuracy could easily be increased to 2-3%. The burning layer of fuel was represented by a flat illuminator with uniformly luminous matt surface. The measurements were made and for each unit of sub-division a table of 64 local values of absorption factor was drawn up. There was no need to make 64 such tables; because of the symmetry of the model only 16 were required. The tables were then used to calculate absorption factors from formulae (3) and the distribution of incident fluxes on the walls of the model was determined for the case of a uniformly radiant medium and a fuel layer. Examples of radiant flux distribution of the model were plotted. Attention has recently been drawn to radiation

Card 5/6

44

87758

8/096/61/000/002/011/014
E194/E155

Optical Modelling of Radiant Heat Exchange in Furnaces

back from the screen tubes. Accordingly the difference between the actual operating conditions of screens and those which are usually assumed was investigated. A study was made of the influence of the degree of blackness of the screen tubes on the heat exchange conditions in furnaces. The method of setting up the model to do this is briefly described and comparative data for tubes with absorption factors of 0.6 and 0.9 shows that alteration of the degree of blackness of the screen tubes has no important influence on the radiant heat exchange in the case considered. Results obtained in tests on the optical model with almost black surface were compared with calculated values for absolutely black tubes and agreement was good. Ways in which the procedure may be further developed are discussed and it is recommended as a useful aid in calculations of heat exchange.

There are 3 figures, 1 table and 11 references: 7 Soviet and 4 English.

ASSOCIATION: Kazakhskiy universitet i TsKTI
(Kazakh University and Central Boiler Turbine Institute) *44*

Card 6/6

3/263/62/000/006/013/015
1008/1208

AUTHORS: Vulis, L.A. and Klinger, V.G.

TITLE: A method of light integration

PERIODICAL: Referativnyy zhurnal, otdel'nyy vypusk. 32. Izmeritel'naya tekhnika, no.6, 1962, 51, abstract 32.6.324.
(Tr. Kazakhk.un-ta, 1960, no.2, 103-108)

TEXT: The possibility of calculating the irradiation of targets, of the radiation of sources if self-absorption is taken into account, and of the radiation barrier, etc., by means of an experimental study of radiation rays on an optical model (the latter serves as a light integrator), is discussed. The simplicity and the sufficient accuracy of the method are illustrated by means of an example, in which the radiation dose received by a disk-like target from a source of particles of the same shape is determined.

[Abstracter's note: Complete translation.]

Card 1/1

KLINGERT, N.V., Eng.

Penstocks

Welded turbine collector. Gidr. stroi. 20 No. 5, 1951.

Monthly List of Russian Accessions, Library of Congress, November 1952. UNCLASSIFIED

KLINEKRT, W.V., inzhener.

Calculation of steel pressure-pipelines. Gidr.stroi. 22 no.10:37-39 0 '53.
(MLRA 6:10)
(Pipe, Steel)

KLINGERT, Nikolay Vasil'evich; KHOKHARIN, Anatoly Kharitonovich;
KHAZANOVA, A.Z. inzh., retsenzent

[Steel pipelines and equalizing reservoirs of hydroelectric power stations] Stal'nye truboprovody i uravnitel'nye rezervuary gidroelektricheskikh stantsii. Moskva, Energiia, 1965. 207 p. (MIRA 1813)

1. Leningradskaya proyektno-konstruktorskaya kontora
"Gidrostal'proyekt"

Klingert, P.V., Inshener.

New developments in the technology of "automats." Stan. 1 instr.
18 no. 4:27 Ap '47. (MLRA 7:11)
(Machinery, Automatic)

1. The action of desoxycholic acid on the adrenal glands of adult rats. A histological and histochemical study. C. I. Parker, J. Ross, L. Louria, and D. K. Elleray. *Cancer*, 1959, **12**, 67-80 (1961). Forty-two adult rats received daily doses of 1.0-3 mg. of desoxycholic acid for 1-10 months. An atrophy of 27-34% of the pituitary glands of the males was observed, but no histological or histochemical changes were found. It is assumed that the depression in the function of the pituitary gland is secondary.

KLINGHER-WASSERMAN, D.

Country : RUMANIA
Category: Human and Animal Physiology. Internal Secretion.
Thyroid Gland

Abs Jour: RZhBiol., No 19, 1958, 88953

Author : Nitescu, I.; Wasserman, L.; Klingher-Wasserman, D.
Govrilito, L.

Inst : Romanian Academy, Iasi Affiliate
Title : The Protective Action of Vitamin A in Dystrophy of
the Thyroid Gland, Caused by Thicuron Derivatives.

Orig Pub: Studii si cercetarii stiint. Acad. RPR Fil. Iasi Mod.,
1958, 7, No 1, 1-16

Abstract: Methylthiouracil was administered in doses of 5 mg/
100 g to rats for a period of 15 days. Typical
changes resulted in the thyroid gland and in the
hypophysis. These changes were less marked (parti-

Card : 1/2

KINCHER-WASSERMANN, D.

Effect of nicotinic acid on the evolution of exanthemas of the rat. I. Nitulescu, L. Wassermann, and D. Kucher. *Arch. exp. Pathol. Physiol.* 1936, Vol. 26, No. 1, 49-61 (1936). The effect of a single administration of small doses (1.5 mg. per g. body weight) of nicotinic acid to animals with a certain exanthema is considered. Nicotinic acid has a favorable effect on the evolution of the months of exanthemas. It does not influence the excretion of the epinephrine. The excretion of epinephrine from the different glands is observed.

KLINGHOFER, L.

Clinical diagnosis of the Kinnelstiel-Wilson syndrome. Orr, betil.
93 no. 15:455-456 13 Apr 1952. (CML 23:3)

1. Doctor, 2. Second Internal Clinic (Director -- Prof. Dr. Gabor
Groniczer), Szeged Medical University.

KLINGHOFFER, L.; SZUTZ, I.

A case of subacute erythroleukemia. Orv. hetil. 94 no.25:684-686 21
June 1953.
(CML 25:1)

1. Doctors. 2. Second Internal Clinic (Director -- Prof. Dr. Gabor
Ozonicszer), Szeged Medical University.

KLINGHOFFER, L.; GABOR, P.

Surgical arteriovenous fistula in the therapy of hypertension. Orv.
hetil. 94 no. 28:776-777 12 July 1953. (CLML 25:1)

1. Doctors. 2. Second Internal Clinic (Director -- Prof. Dr. Gabor
Csontos) and Institute of Surgical Anatomy and Surgery (Director --
Prof. Dr. Gabor Petri) of Szeged Medical University.

"APPROVED FOR RELEASE: 09/18/2001

CIA-RDP86-00513R000723130008-7

KLINGHOFER, L.; ZSOTER, T.

~~Garditis in serum sickness. Orv. hetil. 94 no.45:1252-1253 8 Nov 1953.~~
(CLNL 25:5)

1. Doctors. 2. Second Internal Clinic (Director --- Prof. Dr. György
Oszegicser), Szeged Medical University.

APPROVED FOR RELEASE: 09/18/2001

CIA-RDP86-00513R000723130008-7"

KLINGHOFER, Lasslo, dr.; DAVID, Andras, dr.

Pathological ballistocardiogram. Magy. belorv. arch. 8 no.6:179-183
Dec 55

1. A Gyori Magyei Korhas igazgato: Mike Zoltan dr. II. szam
Belosztalyanak (főorvos: Klinghofer Lasslo dr.) kosleménye.
(BALLISTOCARDIOGRAPHY
in pathol. heart cond. (Hun))

KLINGHOFER, László, dr.; SZABÓ, Rezső, dr.; PINTER, Imre, dr.

Waldenström's purpura hyperglobulinæmica. Orv. hetil. 96 no.4;
107-109 23 Jan 55.

1. A Szegedi Orvostudományi Egyetem II. sz. Belklinikájának
(igazgató: Csonicszter Gábor dr. egészségi tanár) kozleménye.
(HEMORRHAGIC DIAPHERESIS,
thrombopathy)

KLINGHOFER, Lasslo, dr.; DAVID, Andras, dr.

Ballistocardiographic in 50 patients. Orv. hetil. 96 no.8:219-221
20 Feb 55.

1. A Gyori Megyei Korhas (igazgato: Mike Zoltan dr.) II. sz.
Belosztalyamak (foorvost: Klinghofer Lasslo dr.) konkurennya.
(BALYSTOCARDIOGRAPHY,
results in 50 patients)

KLINGHOFFER, Lasslo, dr.,; SLABO, Resso, dr.

Electrocardiographic diagnosis of rheumatic carditis. Orv. hetil.
96 no.29:795-800 17 July 55.

1. A Szegedi Orvostudomanyi Egyetem II. sz. Belklinikajának
(igazgató: Csönicsz Gábor dr. egyet. tanár) koállomány.
(RHEUMATIC HEART DISEASE, diagnosis,
ECG)
(ELECTROCARDIOGRAPHY, in various diseases,
rheum. heart dis.)

KLINGHOFER, L.

EXCERPTA MEDICA Sec.18 Vol.1/4 Cardiovascular Apr 57

933. KLINGHOFER L. and SZABÓ R. 2. Med. Klin., Univ. Szeged; 2. Med. Abt., Komitatskrankenanstalt Györ. Über die Rolle der Elektrokardiographie in der Diagnostik der ventrikulären Hypertrophie an Hand von 100 eigenen Fällen *The role of ECG in the diagnosis of ventricular hypertrophy demonstrated in 100 personal cases* Z. ges. inn. Med. 1956, 11/15 (702—710) Graphs 6 Tables 3

The ECG signs of hypertrophy are described from a differential-diagnostic point of view. Hypertrophy was more frequently diagnosed from the thoracic lead. Standard and thoracic leads, however, frequently supplemented each other. It is demonstrated that in ventricular hypertrophy the form of the ECG curve is not determined by the anatomical conditions.

Florian - Munich

KLINGHAFER, LASZLO, Dr.; MICHAEL BLECHER, Dr.; VILAGI OYULA, dr.; DAVID ANDRAS, Dr.

Ballistocardiography in differentiation of heart complaints of organic
and nervous origin. Magy. belorv. arch. 10 no.2-3;41-42 Apr-June 57.

1. A gybri Magyar Korhaz (igazgato: Mikl Zoltan dr.) II. os. Belosz-
talyanak (foorvos: Klinghafer Laszlo dr.) koslemenyek.
(BALLISTOCARDIOGRAPHY, in various dis,
heart dis., differ. diag. of organic & nerv. heart dis.
(Hun))

KLINGMAN, M.N.; KALINKA, V.D.

Micrometastatic cancers in the spinal cord simulating Duchenne-Aran
spinal amyotropia. Zhur. nevr. i psikh. 61 no.11:1630-1635 '61.
(MIRA 15:2)

1. Klinika nervnykh bolezney (zav. Rafedroy - prof. A.S.Pentsik)
Rishskogo meditsinskogo instituta i 1-ya Rizhskaya gorodskaya
klinicheskaya bol'ница (glavnyy vrach K.F.Bergman).
(SPINAL CORD-CANCER) (ATROPHY, MUSCULUM)
(MUSCLE-CANCER)

KLINGMAN, M.N.

Phenylin treatment of thrombotic cerebral processes. Sov. med.
25 no.4:119-120 Ap '62. (MIRA 15:6)

1. Is kliniki nervnykh bolezney (zav. - prof. A.S. Pentsik)
Riishskogo meditsinskogo instituta i nevrologicheskogo otdeleniya
Riishskoy klinicheskoy bol'nitsy No.1 (glavnyy vrach K.F. Bergman).
(THROMBOSIS) (ENDARTERIUM)

L 25963-66

ACO NR: AP6006421 SOURCE CODE: UR/0317/65/000/011/0060/0061

AUTHOR: Klingner, K. (Engineer, Captain, Member of the National People's army of the GDR)

ORG: None

TITLE: Equipment for military staff vehicles

SOURCE: Tekhnika i vooruzheniya, no. 11, 1965, 60-61

TOPIC TAGS: motor vehicle, military operation, armored vehicle, communication equipment, armed force logistics

ABSTRACT: The military staff equipment carried on motor vehicles in East Germany is briefly described. The leading operation groups usually use armored vehicles of caterpillar or wheel types. The vehicles are equipped with working desks and radio-communication devices of a short-wave range. Telephones are used at stopovers. The main body of the staff personnel is moved in staff vehicles equipped with desks, map boards, cabinets, chairs, typewriters, duplicating and computing machines. In order to increase the working floor space,

Card 1/2

L 25963-66

ACC NR: AP6006421

the lateral walls of the vehicle are made of a collapsible type which, when let down, increases the floor surface 2.5 times. The arrangement of equipment inside the staff vehicle was schematically shown for two versions of which the first one was designed for field operation while the second one was used for rear services. Orig. art. has: 1 figure.

SUB CODE: 15,19 / SUBM DATE: None / ORIG REF: 000 / OTH REF: 000

Card 2/2

KLINCO, V. V., Cand Phys-Math Sci -- (diss) "On the problem of the theoretical determination of the superfine structure of atomic terms." Vil'nyus, 1959. 7 pp (Min of Higher Education USSR. Vil'nyus State U im V. Kapsukas). 150 copies (KL,39-59,101)

8

UGORETS, I.I.; GLAZUNOV, A.A.; SYROMYATNIKOV, I.A.; KASHUNIN, I.S.; POSTNIKOV,
N.A.; RADTSIG, V.A.; UL'YANOV, S.A.; GRUDINSKIY, P.O.; VASIL'YEV, A.A.;
KUVSHINSKIY, M.M.; BAPTIDANOV, L.N.; TARASOV, V.I.; KRIKUNCHIK, A.B.;
SHAPIRO, A.B.; BIBIKOV, V.V.; DVOSHIN, L.I.; KLINEV, I.D.; KARPOV,
M.M.; USPENSKIY, B.S.; CHALIDZE, I.M.; BLOCH, Ya.A.; SHMETKIN, I.S.

Iesif IAkevlevich Gumin; obituary. Elek.sta.26 no.12:58 D 155.
(Gumin, Iesif IAkevlevich, 1890-1955) (MIRA 9:4)

KLINGOF, I.D.

AUTHOR: Lebedev, A.N., Cand.Tech.Sci.
and Klingof, I.D., Engineer SOV/96-58-7-18/22

TITLE: The characteristics, reserves and consumption of fuels produced
in India (Kharakteristika dobivayemykh v Indii topliv, ikh zapasy
i potebleniye.)

PERIODICAL: Teploenergetika, 1958, Vol. 5 No. 7. pp. 85-86 (USSR)

ABSTRACT: The article opens with a brief historical survey of coal mining in
India. Figures are given for 1956 production. The productivity of
labour is very low because mechanisation is slight. The main
properties of coals from a number of fields are described, and the
leading properties of some are tabulated. The properties of
Indian crude oil are given; it is of normal viscosity, low ash
but high sulphur content. Prospective increases in coal and oil
production are considered. There is 1 figure, 1 table and 4
literature references (1 Soviet and 3 English)

1. Fuels - India 2. Fuels - Properties 3. Fuels - Availability
4. Fuels - Consumption

Card 1/1

SOV/96-59-2-15/18

AUTHORS: Lebedev, A.N., Candidate of Technical Sciences
Klingof, I.D., Engineer

TITLE: Power Engineering in India (Energetika Indii)

PERIODICAL: Teploenergetika, 1959, Nr 2, pp 89-91 (USSR)

ABSTRACT: The article opens with a brief review of power engineering developments in India since 1946; curves of installed capacity and power generated in different years being given in Fig 1. Brief descriptions are then given of the power stations at Bokaro, constructed in 1953 and Trombey. There are 3 figures.

Card 1/1

KLINGOFER, H.

"Professional Libraries as Valuable Assistance in the Improvement of Professional Qualifications", p. 74, (IECHNIK, Vol. 27, No. 2, Feb. 1954, Warszawa, Poland)

SG: Monthly List of East European Accessions, (EEL), LC, Vol. 4, No. 5, May 1955, Uncl.

KLINI, St. K.
KLINI, St. K.

"Introduction to Metamathematics," Izdat. Inostr. Lit., Moskva, 1957.
2 526 pp. 32.25 R.

KLINIKA, M. (Praha 12, Nad Petruskou 4.)

An improved snare for the extraction of concrements from the ureter.
Roshl. chir. 37 no.5:332-335 May 58.

(URONTES, dis.

concrections, improved snare for extraction (Cs))

KLINIKOVA, L.A.; TORBOV, V.I.; GORDEYEV, I.V.

Crystallisation of indium phosphide from a gaseous phase.
Izv. AN SSSR. Neorg. mat. 1 no.12:2100-2101. D 1965.

(MRA 1602)

1. Institut novykh khimicheskikh problem AN SSSR. Softaikov
June 29, 1965.

VASIL'YEV, Dmitriy Vasil'yevich, zasl. deyatel' nauki i tekhniki
RSFSR; MIKHAYLOV, Vladimir Aleksandrovich; NORNEVSKIY,
Boris Ivanovich; VYLKOST, V.D., retsenzent; KUTASIN, B.P.,
retsenzent; KLINIMA, Ye.V., red.

[Automation of ship equipment] Avtomatizatsiya sudovykh
ustanovok. Pod red. D.V.Vasil'eva. 2. izd. perer. i dop.
Leningrad, Sudostroenie, 1965. 607 p. (MIRA 19:1)

KLININ, Ye.V., kand.tekhn.nauk, dots.

Explosions and burns during the testing of high-voltage apparatus.
Elektrichesvo no.6:91-92 Je '58. (MIRA 11:6)
(Electric transformers--Testing) (Condensers (Electricity)--Testing)

KLINISKII, Ye.L.

Use of nitroglycerin in evaluating coronary blood circulation.
Kaz. med. zhur. 41 no.3:23-25 My-Je '60. (MIRA 13:9)

1. Iz l-y kafedry terapii (sav. - deystv. chlen AMN SSSR, prof. M.S. VOVSI) Tsentral'nogo instituta usovershenstvovaniya vrachey.
(NITROGLYCERIN—PHYSIOLOGICAL EFFECT)
(CORONARY VESSELS—DIAGNOSIS)
(ELECTROCARDIOGRAPHY)

SHEVCHENKO, M.A.; KLINIYCHUK, Ye.M.; KAS'YANCHUK, R.S.

Purification of water by removing phenols and petroleum products
by ozonisation. Ukr.khim.shur. 3C no.5:527-530 '64.

(MIRA 18:4)

1. Institut obshchey i neorganicheskoy khimii AN UkrSSR.

KLINEK, R.

Effect of ultraviolet rays on the blood sugar level in frogs
during hibernation and artificial rewarming. Acta physiol.polon.
11 no.5/6:773-774 '60.

1. Z Zakladu Fizjologii Pomorskiej A.M. w Szczecinie, Kierownik:
prof.dr E.Mietkiewski.

(HIBERNATION blood)
(BLOOD SUGAR)
(ULTRAVIOLET RAYS)

KLINKE, Romuald

Effect of ultraviolet rays on the blood sugar level in frogs in hibernation and in various environmental temperatures. Roczn. pom. akad. med. Swierczevski. 7:271-282 '61.

1. Z Zakladu Fizjologii Pomorskiej Akademii Medycznej Kierownik: prof. dr Eugeniusz Mietkiewski.

(ULTRAVIOLET RAYS) (HIBERNATION blood)
(BLOOD SUGAR) (TEMPERATURE)

MIETKIEWSKI, Eugeniusz; KLINKA, Romuald

On the effect of ultraviolet rays on the blood sugar level in rabbits. Acta physiol. Pol. 15 no.5:623-634 5-3 '64

I. Z Zakladu Fizjologii Akademii Medycznej w Szczecinie
(Kierownik: prof. dr. E. Mietkiewski).

KLINKE, Ronald; KAMYSZEW, Antoni; FAFROWICZ, Biruta

On the effect of streptomycin and dihydrostreptomycin on chronaxie
of the rabbit ear ~~Micrath.~~ Roczn. Pom. akad. med. Swierczewski
10:217-235 '64.

1. Z Zakladu Fisiologii Pomorskiej Akademii Medycznej (Kierownik:
prof. dr Eugeniusz Mietkiewski) i z Kliniki Ptyzjatrycznej
Pomorskiej Akademii Medycznej (Kierownik: prof. dr Zbigniew
Garnuszewski).

KLINKER, L.

KLINKER, L. Meteorologic effect on fading of short duration which occur in ultra short-wave radio transmission. Tr. from the German. p. 212.

Vol. 60, No. 4, July/Aug. 1956
IDEJAPAS
SCIENCE
Budapest, Hungary

Sov. East European Accession, Vol. 6, No. 2, Feb. 1957

KLINKIEWICZ, F.

POLAND/Chemical Technology - Chemical Products and Their Application, Part 4. - Dyeing and Chemical Treatment of Textile Materials.

H-33

Abs Jour : Ref Zhur - Khimiya, No 7, 1958, 23558
Author : F. Klinkiewicz.
Inst :
Title : Development of Correct Warp Sizing Method Depending on Warp Kind (Combed, Card) and Wool Fiber Content.
Orig Pub : Przem. wlokienniczy, 1956, 10, No 10, Biul. Inst. Wlokienn. 19.

Abstract : A recipe for size preparation for warp with various contents of wool fibers (0, 60 and 100%) was developed. The size consists of potato starch and bone glue with an addition of wax emulsion as softener. The ingredients of the wax emulsion are: synthetic wax, olein, glycerin and ammonium salts. It is recommended also to split the strach.

Card 1/1

KLINKO, M.; OLTVANYI, O.

"The nuclear physics situation in Hungary; Dr. Lajos Janossy and Dr. Sandor Szalay, professors of nuclear physics on the present results and plans."

p. 10 (Ujítók Lapja) Vol. 9, no. 22, Dec. 1957
Budapest, Hungary

SO: Monthly Index of East European Accessions (EEAI) LC. Vol. 7, no. 4,
April 1958

Achievement of the Department of Textile Engineering at Budapest
Polytechnic. p. 45

LEZA PROMISHEGST. Vol. 5, No. 1, 1956

Sofia, Bulgaria

So. East European Accessions List Vol. 5, No. 9 September, 1956

KLINKO, Mark

One of the most modern photograph laboratories in Europe is at the
Hungarian Telegraph Agency. Ujít lap 12 no.4:12-13 25 F '60.

UDOVITSKIY, S.; SHEMETS, A.; LILOV, A. (Chernovtay); KLINKOV, I. (Serpukhov Moskovskoy obl.); TERTYCHNYY, F. (Makeyevka Donetskoy obl.); BOROD'KO, I. (Vorkuta, Komi ASSR); BAZUKIN, P. (Novokuznetsk, Kemerovskoy obl.)

From the editor's mail. Sov. profsoiuzy 20 no.2:32-33 Ja'64.
(MIRA 17:2)

1. Zaveduyushchiy yuridicheskim sektorom Ukrainskogo respublikanskogo soveta professional'nykh soyuzov, Kiyev (for Uдовитский). 2. Konsul'tant yuridicheskogo sektora Ukrainskogo respublikanskogo soveta professional'nykh soyuzov, Kiyev (for Шемет). 3. Neshtatnyy korrespondent zhurnala "Sovetskiye profsoyuzy" (for Бород'ко).

L 52621-65 EWT(1)/EWP(e)/EWT(m)/EWP(l)/I/EWP(t)/EEC(b)-2/EWP(s)/EWA(h)/EWA(c)

Pz-/-Peb/Fi-4 LWP(c) JD/QO/AT

ACCESSION NR: AP5014075

UR/2363/65/001/004/0478/0179

AUTHOR: Grinberg, Ya. Kh.; Medvedeva, Z. S.; Klinkova, L. A.

53

TITLE: Preparation of boron phosphide single crystals

52

SOURCE: AN SSSR. Izvestiya. Neorganicheskiye materialy, v. 1, no. 4, 1965, 478-479

B

TOPIC TAGS: compound semiconductor, boron phosphide, high purity boron phosphide, synthesis, single crystal growth, chemical transport reaction, physical property

ABSTRACT: Synthesis of high-purity (99.998%) microcrystalline boron phosphide powder and a technique of growing boron phosphide single crystals have been developed to produce crystals of the purity and size suitable for measurement of physical characteristics. Difficulties encountered in preparation of this refractory compound semiconductor were emphasized. The purity achieved by the process described was almost an order of magnitude higher than in previous preparations. A chemical transport reaction with iodine vapors was used for growing the single crystals. The reaction involving a diffusion mechanism produced 1-1.5-mm large crystals. Morphology of the crystals was described and x-ray crystallographic data were given. Microhardness of the crystals was measured and found to be somewhat different than

Card 1/2

L 52621-65

ACCESSION NR: AP5014075

previously reported. The average thermoelectric power in the 20-150C range was about 150 $\mu\text{V}/\text{degree}$. All crystals displayed n-type conductivity. Orig. art. has figure. [JK]

ASSOCIATION: Institut obshchey i neorganicheskoy khimii im. N. S. Kurnakova Akademii nauk SSSR (Institute of General and Inorganic Chemistry, Academy of Sciences SSSR)

SUBMITTED: 18 Jan 65

ENCL: 00

SUB CODE: AF, GC

WT REF SERV: 007

OTHER: 009

ATT PRESS: 4010

102
Card 2/2

I. 10256-66 ENT(a)/T/EMP(t)/EMP(b)/EMI(c) IIP(c) JD
ACC NR. AP6001226

SOURCE CODE: UR/0363/65/001/012/2100/2101

29
23

AUTHOR: Klinkova, L. A.; Torbov, V. I.; Gordyev, I. V.

ORG: Institute of New Chemical Problems, Academy of Sciences SSSR (Institut novykh khimicheskikh problem Akademii nauk SSSR)

TITLE: Crystallization of indium phosphide from the vapor phase

SOURCE: AN SSSR. Izvestiya. Neorganicheskiye materialy, v. 1, no. 12, 1965,
2100-2101

TOPIC TAGS: indium phosphide, crystal growing, chemical transport reaction, single crystal, crystallization

ABSTRACT: A preliminary study has been made of the effect of chemical transport reaction conditions on the preparation of InP single crystals from the vapor phase. The experiments were conducted in sealed evacuated (up to 6×10^{-6} mm Hg at 20°C) quartz ampoules using polycrystalline cubic InSb ($a = 5.869 \text{ \AA}$) as the starting material. The transport temperatures were: in the heterogeneous reaction zone, 950°C; in the crystallization zone, 900°C. The transporting agents were I or, for a faster reaction, InI. Depending on the transporting agent, concentration, and ampoule diameter the following InP crystals were prepared: 1) n-type crystals of cubic modification up to 2 mm; 2) dendrites up to 3 mm; or 3) polyhedral crystals up to 2 mm. The prerequisites for controlled growing of InP single crystals are an elucidation of the mechanism of the reaction mixture transport to the crystallization zones, and the

Cord 1/2

UDC: 546.682'181.1.548.19

L-10236-66

ACC NR. AP6001226

relation between the transport process and crystal growth. Orig. art. has: 1 fig.
[BO]

SUB CODE: 20/ SUBM DATE: 29Jun63/ ORIG REF: 001/ OTM REF: 011/ ATD PHMS:
4168

OC
Card 2/2